

****FULL TITLE****

*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***

****NAMES OF EDITORS****

GOLF: a new proxy for solar magnetism

S. Mathur

Indian Institute of Astrophysics, Koramangala, Bangalore 560034, India

Instituto de Astrofísica de Canarias, 38205, La Laguna, Tenerife, Spain

S. J. Jiménez-Reyes

Instituto de Astrofísica de Canarias, 38205, La Laguna, Tenerife, Spain

R. A. García

Laboratoire AIM, CEA/DSM-CNRS-Université Paris Diderot; CEA, IRFU, SAp, Centre de Saclay, F-91191, Gif-sur-Yvette, France

Abstract. Solar magnetism is measured with different indexes: for instance the MPSI and the MWSI, number of sunspots, radio flux at 10.7 cm, Ca II K, Mg II K, EUV, He I or L_{α} . Bachmann & White (1994) had compared these indicators of the solar activity showing a hysteresis of the solar cycle variations and a time lag between these indices not related to instrumental effects. Later on, Özgüç & Ataç (2001) extended this study of hysteresis phenomenon between Flare index and other solar indices (mean magnetic field, coronal index). In its original working configuration, GOLF/SoHO was able to measure during 26 days the solar mean magnetic field (García et al. 1999). We check here if the velocity data could be used as another solar magnetism proxy with the advantage of having a duty cycle $\geq 95\%$ during the last 12 years. We will compare the GOLF data with some of the above-mentioned solar activity indexes.

1. Introduction

GOLF¹ is a resonant scattering spectrophotometer dedicated to the study of low-degree acoustic (e.g. García et al. 2001) and gravity modes (e.g. Mathur et al. 2007; García et al 2008). With a sampling time of 10s, it measures the Doppler velocity on the doublet sodium line, NaD1 and NaD2, by integrating two points of $\sim 25 \text{ m\AA}$ width on the same wing. During the last 12 years, measurements have been alternated between the Blue Wing (BW), the Red one (RW) and back to the Blue one (García et al. 2005). Each observed wing corresponds to a different height in the solar atmosphere (Jiménez-Reyes et al. 2003, 2007). In the presence of a magnetic field the solar Na absorption lines are broaden (Robillot, Bocchia and Denis 1993) and, therefore, the Doppler velocity observed by GOLF will be sensitive to the magnetism at the surface of the Sun.

¹Gabriel et al. 1995

2. Time and spectral analyses and comparison with solar indexes

Two of the usual solar activity indexes are the Magnetic Plage Solar Index (MPSI) and the Mont Wilson Sunspots Index (MWSI). The MPSI is the sum of those pixels where the absolute value of the magnetic field is between 10 and 100 Gauss. Whereas the MWSI is calculated for pixels having absolute values greater than 100 Gauss.

GOLF and MPSI see the same magnetic structures but shifted in time, whereas the magnetic structures in the MWSI are completely different (Fig. 1 (left)). Indeed GOLF integrates the full solar disk and thus takes a mean value which is mainly the weak magnetic field as sunspots only represent a few percent of the solar surface. Thus GOLF is more sensitive to small magnetic structures. Meanwhile, MPSI and GOLF spectra present two peaks corresponding to the 26 days periodicity (rotation period) and its first harmonic (Fig. 1 (right)).




Figure 1. On the left: zoom of the time series of GOLF, MPSI and MWSI. On the right: Power Spectrum Density at low frequency of GOLF and MPSI.

We deepen our study by using two more indexes: sunspot numbers (SS) and the radio flux at 10.7 cm (F10.7). We calculate the correlation rates and time lags between them and GOLF. Table 1 shows that the observation of activity is made at different moments depending on the index studied. GOLF data present a delay from 3 to 5 days compared to the other indexes. Moreover, the correlation rate is higher with MPSI than with MWSI or SS as GOLF integrates solar light and is sensitive to weak magnetic field (see Section 2).

For the different periods of time in GOLF observations, we calculate the time lags and correlation coefficients between GOLF and MPSI (Fig. 2). Using 12 years of data, the correlation has a maximum of 69% for a time lag of 3.57 days, meaning that magnetic structures are seen 3.57 days after by the GOLF instrument. This is probably due to a lack of sensitivity towards the solar limb (García et al. 1998). The decrease of the correlation in the RW is due to the lower sensitivity of this operating mode to the broaden of the line in presence of magnetic field.

Table 1. Time lag and correlation coefficient between GOLF and five solar activity indexes

Index	Time lag (days)	Correlation rate
MPSI	3.57	69%
MWSI	3.62	48%
MPSI+MWSI	3.58	68%
SS	3.45	51%
F10.7	4.95	53%




Figure 2. Cross-Correlation between GOLF and the MPSI for different configuration of observation of GOLF as a function of the time lag in days.

3. Using the wavelet technique

Applying the wavelet² technique to the data, we obtained the Wavelet Power Spectrum (Fig. 3 top and middle) as a function of the period of the wavelet and of time. Regions where the power is more significant (in red and black colours) appear in both spectra and almost at the same moment. The same behaviour is observed with the other solar activity indicators. Fig. 3 (bottom) shows the coherency between the wavelet power spectra of MPSI and GOLF data shifted by 3.57 days. The maximum coherency is observed around a period of 26 days.

4. Conclusions

A 26-days periodicity appears in GOLF data (peak at 26.8 days and first harmonic at 13.4 days in the Power Spectrum). This signal had been seen by Claverie et al. (1982, 1983) and correctly interpreted later by Duvall et al. (1983) and Edmund & Gough (1983). Larger periodicities cannot be found as a

²Wavelet software was provided by C. Torrence and G. Compo and is available at URL: <http://paos.colorado.edu/research/wavelets/>




Figure 3. Wavelet Power Spectrum as a function of time for the MPSI (top panel) and GOLF data (middle panel). Wavelet power spectrum coherency between GOLF and MPSI data (bottom panel).

low-frequency filter is applied to GOLF data. GOLF is sensitive to weak magnetic field as it integrates the full solar disk. However, GOLF observes solar magnetism a few days after other classical activity indexes but with a very high duty cycle ($\geq 95\%$ in 12 years) compared to the MPSI ($\sim 75\%$ during the same period) and with a very fast cadence ($dt=10s$).

The wavelet analysis of the integrated velocity could be a very good tool in asteroseismology to obtain an indication of the full projected orbital velocity and not half of the rotation period as it is the case with the power spectral density.

Acknowledgments. This work has been partially funded by the Spanish grant PENAyA2007-62650 and the CNES/GOLF grant at the SAP-CEA/Sacaly. SOHO is a cooperation between ESA and NASA.

References

- Bachmann, K. T., & White, O. R. 1994, *Sol. Phys.*, 150, 347
- Claverie, A., Isaak, G. R., McLeod, C. P., et al. 1982, *Nature*, 299, 704
- Claverie, A., et al. 1983, *Nature*, 301, 589
- Duvall, Jr., T. L., Jones, H. P., & Harvey, J. W. 1983, *Nature*, 304, 517
- Edmunds, M. G., & Gough, D. O. 1983, *Nature*, 302, 810
- García, R. A., Roca-Cortés, T., Régulo, C. 1998, *A&A*, 128, 389
- García, R. A., Boumier, P., Charra, J., et al. 1999, *A&A*, 346, 626
- García, R. A., Régulo, C., Turck-Chièze, S., et al. 2001, *Solar Phys.*, 200, 361
- García, R. A., Turck-Chièze, S., Boumier, P., et al. 2005, *A&A*, 442, 385
- García, R. A., Jiménez, A., Mathur, S., et al. 2008, *Astronom. Nachrichten*, 329, 476
- Jiménez-Reyes, S. J., García, R. A., Jiménez, A., et al. 2003, *ApJ*, 595, 446
- Jiménez-Reyes, S. J., Chaplin, W. J., Elsworth, Y., et al. 2007, *ApJ*, 654, 1135
- Mathur, S., Turck-Chièze, S., Couvidat, S. & García, R.A. 2007, *ApJ*, 668, 594
- Ozgüç, A., & Ataç, T. 2001, in *IAU Symp.* 203, p125
- Robillot, J.- M., Bocchia, R., & Denis 1993, *Proceedings of the Vth IRIS/GOLF93*